

## Supplementary Material

**Table S1.** Clinical and dermoscopic features.

	Scalp	Back	P value
Patients, n	30	63	
Sex (%)	M 27 (90%) F 3 (10%)	M 39 (62%) F 24 (38%)	0,01
Breslow (mm)	Mean 5,54 (range 0,8-39 mm)	Mean 5,35 (range 0,8-25 mm)	
Ulceration	9 (30%)	32 (51%)	0,09
Satellites	4 (13%)	8 (13%)	0,7*
Demoscopic criteria	N (%)	N (%)	
Milky Red	0	5 (8)	0,1*
Blue-Black	12 (40)	25 (39.7)	1
White lines e crisalidi	10 (34)	37 (59)	0,03
Pseudonetwork	4 (4.4)	0	0,009
Atypical pigment network	6 (20)	25 (39.7)	0,01
Gray-blue areas	14 (47)	20 (32.8)	0,2
Atypical vascular pattern	12 (40)	29 (47.1)	0,7
Streaks	9 (30)	11 (18.5)	0,2
Blotches	14 (47.7)	36 (58.2)	0,4
Irregular dots and globules	7 (24)	18 (38.6)	0,7
Regression areas	9 (30)	14 (22.3)	0,6
Hyper pigmented follicular openings	10 (43.4)	0	<0,001
Pigmented rhomboidal structures	5 (16.7)	0	0,002
Dark blotches and obliterated hair follicles	12 (40)	1 (1.6)	<0,001
Concentric circles	1 (3.4)	1 (1.6)	0,5
Gray dots/clods or Gray circles	5 (16.7)	0	0,003
Absence of pigment network	6 (20)	2 (3.2)	0,01
Arborising blood vessels	2 (6.7)	0	0,1*
Leaf-like areas	0	1 (1.6)	1*
Blue-gray ovoid nests	1 (3.4)	2 (3.2)	1*
Microulceration	3 (10)	4 (6.4)	0,7*
Structureless whitish areas	1 (3.4)	5 (8)	0,6*
Large ulceration	6 (20)	26 (41.3)	0,07
Linear-irregular vessels (serpentine)	2 (6.7)	0	0,1*
Structureless pink areas	3 (10)	4 (6.5)	0,8*
White Collarette	2 (6.7)	1 (1.6)	0,2*
Vascular lacunae	2 (6.7)	0	0,1*
Rainbow pattern	0	1 (1.6)	1*

\*P value calculated using Fisher's exact test.